

Technical Data Sheet

Product	Anti-Hu CD71 Purified	
Cat. Number/Size	11-235-C025	0.025 mg
	11-235-C100	0.1 mg

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Antigen	CD71
Clone	MEM-75
Format	Purified
Reactivity	Human
Application	FC (QC tested), IP, ICC
Application details	Flow cytometry: Recommended dilution: 0.5-4 µg/ml.
Isotype	Mouse IgG1
Specificity	The antibody MEM-75 reacts with an extracellular epitope of CD71 antigen (transferrin receptor), a 95 kDa type II homodimeric transmembrane glycoprotein expressed on activated B and T lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes. The antibody MEM-75 does not block binding of transferrin to the receptor.
Other names	TfR1, TfR, TR, Trfr, T9, p90, TFRC
Workshop	HLDA IV: WS Code A 45; HLDA V: WS Code T T-165
Immunogen	NALM-6 human pre-B cell line
Entrez Gene ID	7037
Gene name	TFRC
NCBI Full Gene Name	transferrin receptor
UniProt ID	P02786

Concentration	1 mg/ml
Preparation	Purified by protein-A affinity chromatography.
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage and handling	Store at 2-8°C. Do not freeze.

Images and References www.exbio.cz

The product is intended For Research Use Only. Diagnostic or therapeutic applications are strictly forbidden. Products shall not be used for resale or transfer to third parties either as a stand-alone product or as a manufacture component of another product without written consent of EXBIO Praha, a.s. EXBIO Praha, a.s. will not be held responsible for patent infringement or any other violations of intellectual property rights that may occur with the use of the products. Orders for all products are accepted subject to the Term and Conditions available at www.exbio.cz. EXBIO, EXBIO Logo, and all other trademarks are property of EXBIO Praha, a.s.